

Kenyon College

Digital Kenyon: Research, Scholarship, and Creative Exchange

Kenyon Summer Science Scholars Program

Summer Student Research Scholarship

Summer 2009

The relationship between body surface area and morphological midgut growth of the tobacco hawkmoth, *Manduca sexta*

Nikolas Tun

Follow this and additional works at: <https://digital.kenyon.edu/summerscienceprogram>



Part of the [Biology Commons](#)

Recommended Citation

Tun, Nikolas, "The relationship between body surface area and morphological midgut growth of the tobacco hawkmoth, *Manduca sexta*" (2009). *Kenyon Summer Science Scholars Program*. Paper 102.
<https://digital.kenyon.edu/summerscienceprogram/102>


This Poster is brought to you for free and open access by the Summer Student Research Scholarship at Digital Kenyon: Research, Scholarship, and Creative Exchange. It has been accepted for inclusion in Kenyon Summer Science Scholars Program by an authorized administrator of Digital Kenyon: Research, Scholarship, and Creative Exchange. For more information, please contact noltj@kenyon.edu.



Image from <http://entomology.unl.edu/k12/caterpillars/hornworm/hornwormpage.html>

The relationship between body surface area and morphological midgut growth of the tobacco hawkmoth, *Manduca sexta*

Nikolas Tun '12 with Dr. Haruhiko Itagaki, Kenyon College Department of Biology, 43022



Abstract

The idea of a universal scaling relationship is one constantly being explored to help gain a better understanding of the functions of many different organisms. We have continued our work in estimating the body surface areas (BSA) of 2nd, 3rd and 4th instar larvae of the sphinx moth, *Manduca sexta*, of various body weights (BW). Three lab-reared *Manduca sexta* were embedded and sectioned using a cryostat. All transverse sections were digitized using polarization optics. The sections for two individuals were mathematically analyzed to produce a parameterization equation of their body surface areas. This information was used to construct a virtual hornworm of 2nd and 3rd instar *Manduca* and a parameterized equation for each organism was calculated. The surface area for each of these three individuals was found to be 0.303 cm² for 2F (2nd instar) and 0.877 cm² for 3AA (3rd instar).

Introduction

- An animal's size influences multiple aspects of its characteristics including its anatomy, physiology and behavior⁷.
- The relationship between body size and metabolic rate is one aspect of size studied in particular, with larger organisms having a lower metabolic rate per gram weight than smaller organisms².
- This relationship between body weight (BW) and metabolic rate (MR) is related by the equation $MR=a(BW)^b$ with b being the experimentally determined exponent².
- The exponent, b , has been found to equal 0.67 according to the surface hypothesis theory(1883)⁶. However, metabolic rates across different species of animals tend to show a value of b closer to .75. This gives rise to the question regarding the existence of a universal scaling relationship between body mass and metabolic rate^{1,4,9,10}.
- A model organism for studying this relationship of metabolic rate and body mass in a single species is the larva of the tobacco hawkmoth, *Manduca sexta*. This larva grows exponentially, increasing in weight 10,000-fold, and passing through 5 instars in only 18 days³ (Fig. 1).
- The ultimate goal of the project is to measure the midgut surface area, which has been found to comprise approximately 9% of the total weight of an actively feeding larva (Gillen, Itagaki pers. observation), but as a pilot study to refine our methods, we began by measuring the overall body surface area in larvae whose metabolic rates have been measured.
- Determining the scaling relationship between body weight and body surface area will help give us a better understanding of interspecific scaling relationships and through this an increased understanding of interspecific scaling relationships⁵.




Figure 1: Growth of *Manduca* larvae. Image from: http://www.staff.uni-Marburg.de/~dolzer/mas_pics.html

Methods

Previous Work

Manduca sexta were reared from eggs (Carolina Biological Supply, NC) on a prepared diet. Throughout the 5 instars of the *Manduca*'s growth period, larvae were starved and then fed red-dyed food to assist in the visibility of their midgut. The larvae were then frozen and embedded with Histoprep and dry ice to freeze them in a vertical position. 4 bristles from a paintbrush were embedded with each larvae to serve as a reference point between sections. A cryostat kept at -20° C was used to make transverse sections of each larva at 50µm thickness which were placed on a gel-coated slide. Each of these sections was digitized using a Zeiss Stemi 2000C stereomicroscope equipped with a Canon digital camera and polarization optics.

Image Analysis

Once these images were taken, they were imported into Image J, an NIH image program. Measurements were taken on a 2nd and 3rd instar *Manduca* between each of the 4 orientation dots and the centroid was found (Fig. 2). A line was drawn between the centroid and the upper left dot, to serve as a reference between sections. Each image was rotated so that the aforementioned reference line was located on the x-axis to serve as the 0° angle. 36 measurements were taken from the centroid to the surface of each section 10° apart.




Figure 2: Section of a 3rd Instar *Manduca sexta* with orientation dots and added image analysis lines.

Mathematical

Each set of radii values was imported into MAPLE, where the complementary mathematical analysis was performed on them. Jim Boston '09 and Dr. Judy Holdener were responsible for the creation of this mathematical protocol. A parametric model of the boundary curve was taken for 14 sections in the 2nd instar and 17 sections in the 3rd instar. A second program in MAPLE was then used to create a parametric model of the *Manduca* body surface for each instar, in addition to finding the surface area. The centroid for each section was used to determine a space curve that served as the “generating curve” for the body surface.

Analysis

References

- ¹Darveau, C.-A.; Suarez, R.K., Andrews, R.D., Hochachka, P.W. (2002) Allometric cascade as a unifying principle of body mass effects on metabolism.
- ²Glazier, D.S. (2005) Beyond the '3/4 power law': variation in the intra- and interspecific scaling of metabolic rate in animals. Biol. Rev. 80:1-52.
- ³Goodman, W.G., Carlson, R.O. and Nelson, K.L. 1985. Analysis of larval and pupal development in the tobacco hornworm (Lepidoptera: Sphingidae), *Manduca sexta*. Ann. Ent. Soc. Am. 78: 70-80.
- ⁴Hochachka, P.W.; Darveau, C.-A.; Andrews, R.D.; Suarez, R.K. (2003) Allometric cascade: a model for resolving body mass effects on metabolism. Comp. Biochem. Physiol. A 134:675-691.
- ⁵Koziowski, J., Weiner, J. 1997. Interspecific allometries are by-products of body size optimization. The American Naturalist. 149: 352-380.
- ⁶Randall, D., Burggren, W. and French, K. Eckert Animal Physiology. 2002. New York: W.H. Freeman and Company. pp. 675-681.
- ⁷Schmidt-Nielsen, K. (1984) Scaling: Why is animals size so important? Cambridge Univ. Press, New York.
- ⁸Shine, R., Harlow, P.S., Keogh, J.S., Boeadi. 1998. The allometry of life-history traits: insights from a study of giant snakes (*Python reticulatus*). J. Zool. 244: 405-414.
- ⁹Suarez, R.D.; Darveau, C.-A. (2005) Multilevel-regulation and metabolic scaling. J. Exp. Biol. 208: 1627-1634.
- ¹⁰Suarez, R.D.; Darveau, C.-A.; Childress, J. (2004) Metabolic scaling: a many-splendoured thing. Comp. Biochem. Physiol. B 139:531-541.

Acknowledgements

I would like to thank my mentor Dr. Haruhiko Itagaki for his assistance and constant guidance with this project. I would also like to thank Dr. Judy Holdener, Jim Boston '09, and Anna Frutiger '09 for their assistance with the project. Funding for this project was provided by the Kenyon College Summer Science Scholars Program.

Results

- A 2nd, 3rd, and 4th instar *Manduca sexta* were sectioned and digitized.
- The 2nd and 3rd instar *Manduca sexta* were mathematically analyzed using NIH image and MAPLE software programs. A parameterized equation for each of the individuals was found.
- The general equation for each cross section was found to be:

$$a_0 + \sum_{m=1}^n (a_m \sin(m\theta) + b_m \cos(m\theta))$$

- The parametric models of the surface area were constructed so that the data fit the sinusoidal curve of the form:

$$\langle X(u), Y(u) \rangle = \left\langle X_0 + \sum_{k=1}^n (a_k \sin(ku) + b_k \cos(ku)), Y_0 + \sum_{k=1}^n (c_k \sin(ku) + d_k \cos(ku)) \right\rangle$$

- This equation and the equations for the surface areas of the individual sections were used to find a full parametric virtual model of the 2nd (2F) and 3rd (3AA) *Manduca sexta* (Fig. 3).
- From these parameterized models, the surface area for each of the three organisms was calculated to be 0.303 cm² for the 2nd instar (2F), and 0.877 cm² for the 3rd instar (3AA).




Figure 3: Fully modeled 3rd (3AA) and 2nd (2F) instar *Manduca sexta*. Measurements taken in centimeters. Models were created in MAPLE.

Discussion

This research is the continued work of what Anna Frutiger '09 had previously worked on for two years. Methods to determine the body surface area still needs to be perfected further to reduce error on these parameterizations. Despite this, however, we are hopeful that with more data, we will later be able to observe the changes in the midgut in comparison to the changes in the surface area. Although there is currently not enough data at this point in time to conclude as to whether the surface area growth of the *Manduca sexta* tobacco hornworms is allometric or isometric, it is an excellent ongoing project that we hope will provide answers to the question of whether or not a universal scaling relationship exists.